

Anaphylaxis Emergency Action Plan

Patient Name:				Age:	
Allergies:					
Asthma Yes (high risk	for severe rea	nction) [☐ No		
Additional health problem	s besides ana	phylaxis:			
Concurrent medications:					
			of Anaphylaxis		
			elling of lips and/or tongue		
			ess/closure, hoarsene redness, swelling	ess essential establishment of the second establishment of	
GU ⁻			rhea, cramps		
LUNG* shortness of breath, of					
	ART* w	eak pulse, d	izziness, passing out		
Only a few s			Severity of symptoms life-threatening. ACT		
Emergency Action S 1. Inject epinephrine in thigh					
			Auvi-Q (0.15 mg)	☐ Auvi-Q (0.3 mg)	
			EpiPen Jr (0.15 mg)	☐ EpiPen (0.3 mg)	
			nephrine Injection, US (0.15 mg)	SP Auto-injector- authorized gener (0.3 mg)	
			Other (0.15 mg)	Other (0.3 mg)	
Specify others:					
IMPORTANT: ASTHMA IN	HALERS AND/	OR ANTIHIS	TAMINES CAN'T BE I	DEPENDED ON IN ANAPHYLAXIS.	
2. Call 911 or rescue squa	d (before calli	ng contact)			
3. Emergency contact #1: home			_work	cell	
Emergency contact #2: home			_work	cell	
Emergency contact #3:	home		_work	cell	
Comments:					
Doctor's Signature/Date/Pho	one Number				
Parent's Signature (for indiv	/iduals under a	age 18 yrs)/[)ate		

This information is for general purposes and is not intended to replace the advice of a qualified health professional. For more information, visit www.aaaai.org. © 2017 American Academy of Allergy, Asthma & Immunology 4/2017