

Asthma Action Plan

| Green Zone: Well | Give these medic | Give these medicines every day: | | | |
|---|---|--|---|---------------------------|--|
| No signs of asthmaAble to do normal activitiesNo problems while sleeping | MEDICINE: | HOW | мисн: | WHEN: | |
| Peak flow above: (above 80% of best) | | | | | |
| *Rinse mouth after this medicine | | | | | |
| Yellow Zone: Watch Out! | First — give: | | | | |
| Early Signs of Asthma: • Cold symptoms | ■ Albuterol | 2-4 puffs or 1 | nebulizer | 1-3 times in first hour | |
| Coughing day or night Wheezing day or night | ■ Call your Doc | ■ Call your Doctor or Nurse if not in Green Zone after first hour. Next — if asthma is better after first hour, you may give: | | | |
| Funny feeling in chest | Next — if asthma | | | | |
| My first sign: | ■ Albuterol | 2-4 puffs or 1 | nebulizer | every 4 hours as needed | |
| • Peak flow: | Call your Doctor or Nurse if: Albuterol needed more often than every 4 hours. Albuterol needed every 4 hours for more than 1 day. Keep taking other Green Zone medicines. | | | | |
| Red Zone: EMERGENCY! | First — give <i>now</i> : ■ Albuterol | 6 puffs or 1 | nehulizer | | |
| Late Signs of Asthma:Tight chestBreathing hard or fast | | ■ AND call your Doctor or Nurse. | | | |
| Using neck or stomach muscles to breathe Constant coughing Trouble talking or walking | | - | t reach your Doctor or Nurse <i>immediately,</i> give: 6 puffs or 1 nebulizer | | |
| VomitingLips or nails blue | oral steroid) ■ AND go to the nearest emergency room or call 911. | | | | |
| Peak flow below: (below 50% of best) | | | | | |
| | | | Phone nu | umber of Doctor or Nurse: | |
| Patient/Parent/Guardian Signature | | Date | | | |
| RN/MD Signature | | Date | Night: | | |